

WP3: Verification methods for terrestrial CO2 sources and sinks and carbon stocks

Responsibles:

P. Smith & M. Kuhnert (UNIABDN)

P. Peylin & M. McGrath (CEA-LSCE)

April, 2021


Teleconference



This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 776810

WP3 “Pre-operational system”


Pre-operational system for CO₂ fluxes from terrestrial ecosystems



WP3 “View from within”

Pre-operational system for CO₂ fluxes from terrestrial ecosystems

T3.2





Toward operationalisation

Modeling system consolidation

- Preparation of forcing
- Meteo forcing
 - Land cover
 - Atmospheric CO2

- Running Bottom-up models
- process-based
 - data-driven

- Running Top-down models
- CSR
 - CIF

New estimates up to year -1 available end of September

Jan

T3.1

June

T3.2

July

T3.3

Sep.

T3.4

⇒ New test this summer 2021



WP3 – MAIN ACHIEVEMENT M19-M36

- **T3.1: UPDATE state-of-the-art driving datasets**
- T3.2: UPDATE the bottom-up budget of terrestrial CO₂ fluxes using a few complementary models (statistical or process-based).
- T3.3: UPDATE the Europe-wide inversions of NEE using in situ data of atmospheric CO₂ and a high-resolution transport model.
- T3.4: Develop new multi-data model fusion strategies to investigate CO₂ budgets and trends; Specific focus on Eastern Europe.



T3.1 Collate state-of-the-art driving data

T3.1 Input data sets

Input for
models

Evaluation
data

GHG
budget
component

Data set	Name/ model	Inst.	Coverage	Resoluti on	Time frame	Contact in the project
Land use	HILDA ³	KIT	global	1 km	1900-2010	Richard Fuchs ^a , Karina Winkler ^b
Biomass	HILDA ³	WU	global	1 km	1900-2010	Martin Herold ^c
N-Deposition	EMEP model ⁴	JRC		1 km	2010	Frank Dentener ^d
Erosion	PESERA ⁵	ESDAC	Europe	1 km	2004	Emanuel Lugato ^e
Soil data	LUCAS ⁶	ESDAC	Europe	1 km/10 km	2015	Emanuel Lugato ^e
Climate	UERRA ⁸	ECMWF	Europe	11 km	1961-2017	Richard Engelen ^f
Flux data	FLUXNET network		global	sites	diverse	Dario Papale, Werner Kutsch
Fertiliser application rates	CAPRI	JRC	Europe	0.25°	2000-2012	Adrain Leip ^g
Crop management timing		UNIABD N	EU28	0.25°	2000-2015	Matthias Kuhnert ^h
Fresh water fluxes		ULB	Europe	0.1°	2016	Ronny Lauerwald ⁱ
Ocean coastal fluxes		UiB	Northern Europe	0.125°	1997-2016	Are Olsen ^j
Forest management		WU	Europe	0.125°	2000-2015	Mart-Jan Schelhas ^k , Gert-Jan Nabuurs ^l
Grassland management		CEA-LSCE	Europe	0.5°	1860-2012	Philippe Ciais ^m
atmospheric CO2	FOCAL ¹⁰	UBremen	global	2 km	2015-2016	Maximilian Reuter ⁿ



T3.1 Collate state-of-the-art driving data

Climate data

Land use/cover
data

Soil
properties/
soil erosion

Cropland
management

Grassland
management

Forest
management

N-Deposition
data

River fluxes

Coastal fluxes/
ocean fluxes

Fluxdata

XCO₂ from
OCO-2



Ongoing work



Collection complete




Different data access

T3.1 Collate state-of-the-art driving data


Management data for different land use were collected and provided

Cropland management
management




Nishina et al., 2016

Grassland management



Chang et al., 2016


Forest



Schelhaas et al., 2018

Coastal ocean fluxes

- ❖ Based on surface ocean pCO₂ measurements: data driven approach to map observed pCO₂ data with a set of predictors (e.g. SST, chlorophyl, Mixed layer depth)
 - ❖ Different statistical models (Random Forest (Becker et al, in prep), Multi-Linear Regression (Becker et al, 2021))
 - ❖ 1998 - 2019, monthly pCO₂, daily air-sea fluxes; 0.125° resolution
- ⇒ **CO₂ fluxes estimated from pCO₂**




Coastal ocean fluxes

- ❖ Merging the coastal ocean fluxes with open ocean fluxes
- ❖ Choosing the coastal product where it is available
- ❖ Combined product covering 2020 will be available in early August



Coastal fluxes, RF

(Becker et al., in prep)




Open ocean fluxes

(Rödenbeck et al., 2014)



Combined product



Average fluxes for 2019

Lateral C fluxes and associated CO₂ emissions from inland waters

H. Zhang, C. Gommet, G. Laruelle, R. Lauerwald, P. Ciais and P. Regnier (ULB, Belgium & IPSL-LSCE, France)

CO₂ evasion from European inland waters using an empirical model (based on GLORICH observational data)

0.1° x 0.1° climatology (mean over the last 3 decades)

Key Limitations

- No temporality (seasonal & IAV)
- No linkage with terrestrial C leaching (DOC + POC + CO₂)




Petrescu et al., ESSD, 2021


Lateral redistribution of carbon in Europe using a land surface model (ORCHIDEE-C_{lateral})

- Numerically efficient erosion scheme based on MUSLE
- High spatial-resolution scheme (3')
- « Smart » upscaling strategy (0.5°)

Erosion scheme in ORCHIDEE-C_{lateral}



ORCHIDEE-C_{lateral}



Zhang et al., JAIMEs, 2020





- Terrestrial fluxes
- Soil erosion
- Soil DOC, CO₂ leaching
- Riverine transport
- POC sedimentation & OC decomposition
- River CO₂ outgassing
- **Impact of lateral C fluxes on soil C stocks**

Zhang et al., in prep.



Lateral fluxes



Lateral fluxes with ORCHIDEE-C_{lateral}



12 catchments (time)


57 catchments (space)

Zhang et al., in prep.



River CO₂ evasion with ORCHIDEE-C_{lateral}

Mean flux from observed climatology




Seasonal variability and IAV from ORCHIDEE-C_{lateral}




Lake CO₂ evasion from observations

Seasonal variability from upgraded empirical model

Multiple Linear Regressions to predict pCO₂ from predictors (Temperature, Surface, Altitude, Latitude)




Seasonally-resolved inland water CO₂ evasion from Europe



Laruelle et al., in prep.

To conclude: European land-river C budget with lateral fluxes

(c) DOC




Lateral C loss: 47 (38+7) Tg C yr⁻¹

Export to sea: 21 Tg C yr⁻¹

CO₂ evasion: 19.5 Tg C yr⁻¹

IW sequestration: 6.5 Tg C yr⁻¹


(b) POC



Next steps

- IAV of lateral C fluxes
- Impact of lateral C fluxes on terrestrial C sink
- Impact of lateral C fluxes on coastal ocean (TBD)

(d) CO₂



Zhang et al., in prep.



WP3 – MAIN ACHIEVEMENT M19-M36

- T3.1: UPDATE state-of-the-art driving datasets
- **T3.2: UPDATE the bottom-up budget of terrestrial CO2 fluxes using a few complementary models (statistical or process-based).**
- T3.3: UPDATE the Europe-wide inversions of NEE using in situ data of atmospheric CO2 and a high-resolution transport model.
- T3.4: Develop new multi-data model fusion strategies to investigate CO2 budgets and trends; Specific focus on Eastern Europe.



T3.2 Bottom up budget of CO2 fluxes

APPROACHES



ORCHIDEE

FluxCom

BLUE

G4M *EFISCEN-space* *EFISCEN*
CBM

EPIC *DAYCENT*

ECOSSE



Compartment



T3.2 Bottom up budget of CO2 fluxes

APPROACHES

Process-based models

Data - driven Models

Bookkeeping approach

Ecosystems

ALL

Forest

Crop

Grass

➤ Benefit from all approaches

➤ **Open to more land surface model (from TRENDY experiment)**

➤ To estimate model uncertainties



All

Vegetation

Soil




Compartment

LULUCF - FRANCE





- EFISCEN Space model initialised on NFI plot data for 12 countries
- Individual tree
- Climate sensitive growth functions
- Harvest patterns derived from repeated observations on NFI plots, for different management intensity classes
- Providing NEP, NBP and harvest flux

Fully new approach for bottom up medium resolution forest carbon flux





Harvest intensity (Nabuurs et al. 2019)


Growing stock




Harvest rate



Germany initial growing stock



Germany harvest rate






EFISCEN Space (WUR)

Growth functions in EFISCEN Space work well

... and result in increment rates close to what countries report



Next steps:

- Include more countries
- Check of results by countries
- Soil carbon module
- Cost module
- Dynamic mortality

VERIFY Input data:

Meteo data

HARMONIE daily dataset: cruhar_v3_1 and shortwave solar radiation (cruhar_v3_2) for 2002-2018;


ERA dataset (cruera_v1.1) for 2019

Land cover


Annual maps from HILDA+ for 2001-2015

IIASA's G4M/FLAM output variables of carbon cycle

Variable	Unit	Description	Temporal aggregation
FCO2_NPP_FOR_FOR	kg C m ⁻² yr ⁻¹	FCO2_NPP_FOR / Fluxes per square meter of forest	Annual
AGB	kg C m ⁻²	Above Ground Biomass	Annual





FCO2_NPP_FOR (kg C m⁻² yr⁻¹) in 2019




AGB (kg C m⁻²) in 2019


What drives the change in NBP trend between 1980-1998 and 1998-2012 in Europe?




T3.2 Analysis of the drivers of CO2 fluxes



T3.2 Analysis of the drivers of CO2 fluxes




T3.2 Analysis of the drivers of CO2 fluxes




NEE driver: GPP or TER?

Pixels classified according to mean temperature and precipitation from 01January1997 to 31December2016




Pixels classified according to mean temperature and precipitation from 01January1997 to 31December2016




NEE driver: GPP or TER?

Fluxcom RS only ANN GPP vs. CSR (valid100km) NEE
 01-January-2006 to 31-December-2015
 Removed a linear fit for each month from the timeseries



Fluxcom RS only ANN GPP vs. CSR (valid100km) NEE
 01-January-2006 to 31-December-2015
 Removed a linear fit for each month from the timeseries





WP3 – MAIN ACHIEVEMENT M19-M36

- T3.1: UPDATE state-of-the-art driving datasets
- T3.2: UPDATE the bottom-up budget of terrestrial CO₂ fluxes using a few complementary models (statistical or process-based).
- **T3.3: UPDATE the Europe-wide inversions of NEE using in situ data of atmospheric CO₂ and a high-resolution transport model.**
- T3.4: Develop new multi-data model fusion strategies to investigate CO₂ budgets and trends; Specific focus on Eastern Europe.





T3.3 Inversion of NEE

- ❖ Based on one system: Jena CarboScope-Regional
- ❖ Two step scheme:
Global inversion (TM3) to provide the lateral transport
Regional EU inversion (STILT) at 0.25° resolution - 1h resolution
- ❖ Fossil fuel emissions and Ocean fluxes are prescribed
- ❖ Solve for terrestrial NEE fluxes for period 2006 - 2019 but using corrections at 0.5° - 3h resolutions
- ❖ Test different land NEE priors and ocean priors
B0: VPRM : simple diagnostic model
B1: Fluxcom : data-driven model
B2: SIBCASA: process-based model
- ❖ Test also different [CO₂] station networks


CarboScope-Regional inversions

Different prior flux models, similar posterior fluxes




T3.3 Inversion of NEE

CarboScope-Regional NEE and Temp. anomalies



T3.3 Inversion of NEE

CarboScope-Regional NEE and Temp. anomalies





T3.3 Inversion of NEE

Key messages

- Posterior NEE strongly driven by atm. CO₂ data (not prior, not ocean)
- But posterior fluxes depend critically on station network; much more than on the prior NEE fluxes
- Interannual variations of Posterior NEE resembles that of prior from VPRM
- CO₂ sink for 2019 was stronger than in 2018 over all Europe, specifically for regions affected by the 2018 drought (north Europe)

Outlook

- Include prior fluxes also from Orchidee
- Include “coastal” ocean flux estimates and lake fluxes
- Additional transport model: footprints from Flexpart / LUMIA system (Guillaume Monteil, U. Lund)
- Use Community Inversion Framework (CIF) to assess transport model errors
- 2020 NEE estimates by Sep. 2021




WP3 – MAIN ACHIEVEMENT M19-M36

- T3.1: UPDATE state-of-the-art driving datasets
- T3.2: UPDATE the bottom-up budget of terrestrial CO₂ fluxes using a few complementary models (statistical or process-based).
- T3.3: UPDATE the Europe-wide inversions of NEE using in situ data of atmospheric CO₂ and a high-resolution transport model.
- **T3.4: Develop new multi-data model fusion strategies to investigate CO₂ budgets and trends; Specific focus on Eastern Europe.**

- Carbon cycle data assimilation systems (**CCDAS**)

⇒ Simulations for late 2021



BETHY-CCDAS

Assimilation of

- Satellite soil moisture
- Satellite L-VOD
- In situ atm. CO2 data
- Nighlights (for FF model)

Two-step assimilation

- Fossil Fuel at 0.1 in FFDAS
- Biogenic fluxes at 0.25 in CCDAS

Optimisation of

around 70 parameters in CCDAS and 1.5M in FFDAS

ORCHIDEE-CCDAS

Assimilation of

- FluxNet in situ data
- Satellite Fluorescence
- In situ atm. CO2 data (using LMDZ model)

Optimisation of

- around 150 parameters
- stepwise assimilation of each datastream

CTDAS

Dynamic a-priori fluxes

Using near real-time statistical data (EU, Eurostat,...) and weather data to fill in short-term variations in fluxes at ~2km

Assimilation of

- ICOS CO2 mole fractions
- $\Delta 14C$, CO, NO2

Optimisation of

Around ~10 parameters per country, sectorial FF emissions, NEE per PFT




BETHY CCFFDAS: Carbon Cycle Fossil Fuel Data Assimilation System

2-step optimization scheme:

1. FFDAS provides fossil fuel emission estimates consistent with observations of nightlights and emissions from point sources.
2. optimized fossil fuel emissions are used in CCDAS to provide terrestrial carbon fluxes consistent with observations from the SMOS satellite as well as atmospheric CO₂ concentrations (**currently running**).


Results from step 2 based on CDIAC fossil fuel emissions (Boden et al, 2016) focussing on Europe (TRANSCOM region):

Mean annual NEP 2010-2015 (+ sink)



Annual average 2010-15 in GtC/yr




Experiment	Europ. flux estimate	
	NEP	NPP
Prior	-0.024±1.19	4.60±0.81
CO ₂	0.18±0.12	4.82±0.16
CO ₂ +SMOS	0.30±0.08	4.82±0.12



Scholze et al., GRL, 2019

CT Europe Lagrangian: joint FF-NEE optimization using radiocarbon and CO₂ mole fractions

PhD research Auke van der Woude






Data assimilation of CO₂ and radiocarbon ($\Delta^{14}\text{C}$) to constrain the sources and sinks of CO₂ in Western Europe.

-> Simulations of hourly varying FF emissions (households, traffic, industry, power production, renewables).


-> Simulations of hourly photosynthesis and respiration by biosphere (2x2km)

-> Simulations of atmospheric transport (CO₂, $\Delta^{14}\text{C}$, CO, NO_x) and weather (renewables and biosphere)



assimilation of $\Delta^{14}\text{C}$ helps separate local sources (**fossil**) from diffuse sinks (**bio**)


CT Europe Lagrangian: joint FF-NEE optimization using radiocarbon and CO₂ mole fractions



- Addition of $\Delta^{14}\text{C}$ (18 samples/month/site) reduces uncertainty of FF *and* NEE
- Estimates come (much) closer to truth for ICOS well-covered countries
- RINGO (H2020 project) network also tested: going closer to FF sources and targeted sampling can help, but not as much as increasing # of sites for CO₂

T3.4 HILDA+ Land Use Change for Europe


HILDA+ land use/cover (LUC) change 1x1km/ global/ 1960-2019 (annual)



HILDA+ for Europe at a glance:

- Six main LUC classes
- Full land transitions = gross change
- Forest classes on top, based on ESA-CCI LC

⇒ **Synthesis LUC dataset with unprecedented spatial, temporal and thematic detail.**



Publication: Winkler et al. (2021): *Land use changes are four times greater than previously estimated*. Nat Comms (in press).

Dataset available at: <https://doi.org/10.1594/PANGAEA.921846>


T3.4 The Eastern European carbon sink

K. Winkler, H. Yang, R. Ganzenmüller, R. Fuchs, F. Chevallier, J. Pongratz and P. Ciais

research ongoing

Carbon sink potential of Eastern Europe

Carbon sink estimates for European regions



Most of carbon sink occurs in Eastern European (EUR-East + European Russia):

- L-VOD AGB (2010-2019) ~**0.46** GtC a⁻¹
- SURF, GOSAT, OCO2 Inversions ~**0.45** GtC a⁻¹
- EFISCEN, CBM and UNFCCC ~**0.32** GtC a⁻¹




T3.4 The Eastern European carbon sink

K. Winkler, H. Yang, R. Ganzenmüller, R. Fuchs, F. Chevallier, J. Pongratz and P. Ciais


research ongoing

Carbon emissions from land use change




ABC trend 2003-10 (Reuter et al., 2017)

Aggregated carbon emissions from BLUE (based on HILDA+) 2010-2019



Eastern Europe



BLUE (HILDA+) $\sim 0.03 \text{ GtC a}^{-1}$ (net emissions)
 → **Forestry and agriculture decrease carbon sink effect from land abandonment.**



Thank you for your attention.



This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 776810



WP3 PARTNERS

Two co – leaders:

University of Aberdeen (**UNIABDN**),
Commissariat à l'énergie atomique (**CEA**),

18 partners:

Joint Research Centre (**JRC**), Karlsruhe Institute of Technology (**KIT**),
Max Planck Society for the Advancement of Science (**MPG**),
Norwegian Institute for Air Research (**NILU**),
International Institute for Applied Systems Analysis (**IIASA**),
Integrated Carbon Observation System (**ICOS**) ERIC,
European Centre for Medium-Range Weather Forecasts (**ECMWF**),
Wageningen Environmental Research (**WEnR**),
University of Bremen (**UBremen**), University of East Anglia (**UEA**),
Vrije Universiteit Amsterdam (**VU**), Lund University (**ULUND**),
University of Bergen (**UiB**), Wageningen University & Research (**WUR**),
Université libre de Bruxelles (**ULB**), Saint Petersburg State University (**SPbU**),
Ural Federal University (**UrFU**), Ludwig-Maximilian Munich University (**LMU**)




Additional slides



T3.2 Bottom up terrestrial CO2 fluxes FLUXCOM data – driven model (MPI)


NEE May 2015



NEE estimates for 2019

VPRM
prior

Fluxcom
prior






T3.3 Inversion of NEE

Inversion using satellite OCO-2 data (LMDZ model)

Annual CO₂ vegetation flux estimated by the 68% uncertainty envelope of the latest two CAMS inversions (blue for the surface-driven one and pink for the satellite-driven one) for the 27-member European Union and United Kingdom together. Positive values indicate a source of CO₂ to the atmosphere.






WP3 – STATUS OF DELIVERABLES M19 - M36


D3.5	Second Bottom-up model simulations	CEA	31/12/2020	Submitted
D3.10	Community Inversion Framework (CIF)	CEA	31/1/2021	Submitted
D3.15	Ground based measurements of column CO2 in Russia and improved estimates of XCO2 from satellite data	KIT	31/7/2020	Submitted

OCO-2 satellite derived Column integrated CO2 conc.

Ex.: August 2018



Evaluation against TCCON data




⇒ Next step: Evaluation against campaign data in St Petersburg

T3.2 EPIC crop model




VERIFY meteo data (CRUHAR v.3.1, 1981-2019)
 VERIFY soil data
 VERIFY land cover



Pan-European gridded model EPIC-IIASA


Variable	Unit	Description	Temporal aggregation
mrso	kg m ⁻² month ⁻¹	Total Soil Moisture Content	1 st day of month
mrro	kg m ⁻² month ⁻¹	Total Runoff, Drainage, and Subsurface Runoff	Monthly sum of daily fluxes
evapotrans	kg m ⁻² month ⁻¹	Total Evapo-Transpiration	Monthly sum
cVeg	kg C m ⁻² month ⁻¹	Carbon in Vegetation	1 st day of month
clitter	kg C m ⁻² month ⁻¹	Carbon in Litter	Monthly sum
cSoil	kg C m ⁻² month ⁻¹	Carbon in Soil	1 st day of month
npp	kg C m ⁻² month ⁻¹	Net Primary Production - total	Monthly sum
npp(a)	kg C m ⁻² month ⁻¹	Net Primary Production – aboveground	Monthly sum
npp(b)	kg C m ⁻² month ⁻¹	Net Primary Production – belowground	Monthly sum
rh	kg C m ⁻² month ⁻¹	Heterotrophic Respiration	Monthly sum
fHarvest	kg C m ⁻² year ⁻¹	C Flux to Atmosphere from harvested crop biomass consumption	Last day of year
yoc	kg C m ⁻² month ⁻¹	soil carbon loss with sediments (water erosion)	Monthly sum
clch	kg C m ⁻² month ⁻¹	leached soluble carbon	Monthly sum

Monthly C stock and fluxes:




Regidded to VERIFY


T3.2 FLUXCOM data – driven model (MPI)




In-situ eddy-covariance carbon fluxes



machine learning



global gridded data sets of predictors





Global gridded estimates of carbon fluxes

*NEW and work
in progress*



- higher spatial resolution & subdaily
- extended and improved training and feature data sets
- new machine-learning methods

Drought effects (2018, 2019)



T3.2 Bookkeeping : land use flux model (BLUE)

- Implemented HILDA+ in BLUE
- Revised model setup for runs at 0.01° degrees (previously 0.25°)
- Total LULC emission estimates are comparable to runs based on LUH2 in size and trends for overall Europe but national differences exist
- Elimination of extreme values (i.e. 1960, 2015) in the BLUE runs based on HILDA+ suggests a higher reliability
- High resolution runs based on HILDA+ capture heterogeneity better and provide a more detailed picture of local sources and sinks



T3.2 Analysis of the drivers of CO2 fluxes

CROPLAND - SPAIN

